

Nucleus References

- Bernoux M, Timmers T, Jauneau A, Brière C, de Wit PJ, Marco Y, Deslandes L. 2008. RD19, an Arabidopsis cysteine protease required for RRS1-R-mediated resistance, is relocalized to the nucleus by the *Ralstonia solanacearum* PopP2 effector. [Plant Cell 20: 2252-64](#).
- Carvalho CM, Santos AA, Pires SR, Rocha CS, Saraiva DI, Machado JP, Mattos EC, Fietto LG, Fontes EP. 2008. Regulated nuclear trafficking of rpL10A mediated by NIK1 represents a defense strategy of plant cells against virus. [PLoS Pathog. 4: Dec 26](#).
- Cheng YT, Germain H, Wiermer M, Bi D, Xu F, Garcia AV, Wirthmueller L, Després C, Parker JE, Zhang Y, Li X. 2009. Nuclear pore complex component MOS7/Nup88 is required for innate immunity and nuclear accumulation of defense regulators in Arabidopsis. [Plant Cell 21; 2503-16](#).
- Deslandes L, Rivas S. 2011. The plant cell nucleus: a true arena for the fight between plants and pathogens. [Plant Signal Behav. 2011 Jan;6\(1\):42-8](#).
- Garcia AV, Blanvillain-Baufumé S, Huibers RP, Wiermer M, Li G, Gobbato E, Rietz S, Parker JE. 2010. Balanced nuclear and cytoplasmic activities of EDS1 are required for complete plant innate immune response. [PLoS Pathog July 1; 6](#).
- Kang HG, Woo Choi H, von Einem S, Manosalva P, Ehlers K, Liu PP, Buxa SV, Moreau M, Mang HG, Kachroo P, Kogel KH, Klessig DF. 2012. CRT1 is a nuclear-translocated MORC endonuclease that participates in multiple levels of plant immunity. [Nat Commun Dec 18;3:1297. doi: 10.1038/ncomms2279](#).
- Kim SH, Kwon SI, Bhattacharjee S, Gassmann W. Regulation of defense gene expression by Arabidopsis SRFR1. [Plant Signal Behav. 4; 149-50](#).
- Lai Z, Li Y, Wang F, Cheng Y, Fan B, Yu JQ, Chen Z. 2011. Arabidopsis Sigma Factor Binding Proteins Are Activators of the WRKY33 Transcription Factor in Plant Defense. [Plant Cell. Oct 11](#).
- Monaghan J, Xu F, Gao M, Zhao Q, Palma K, Long C, Chen S, Zhang Y, Li X. 2009. Two Prp19-like U-box proteins in the MOS4-associated complex play redundant roles in plant innate immunity. [PLoS Pathog. 5, Jul 24](#).
- Noël LD, Cagna G, Stuttmann J, Wirthmuller L, Betsuyaku S, Witte CP, Bhat R, Pochan N, Colby T, Parker JE. 2007. Interaction between SGT1 and cytosolic/nuclear HSC70 chaperones regulates Arabidopsis immune responses. [Plant Cell 19; 4061-76](#).
- Pan H, Liu S, Tang D. 2011. HPR1, a component of the THO/TREX complex, plays an important role in disease resistance and senescence in Arabidopsis. [Plant J. Oct 29. doi: 10.1111/j.1365-313X.2011.04835.x](#).
- Pan H, Liu S, Tang D. 2012. The THO/TREX complex functions in disease resistance in Arabidopsis. [Plant Signal Behav. March 1;7\(3\)](#).

Petersen K, Qiu JL, Lütje J, Fiil BK, Hansen S, Mundy J, Petersen M. 2010. Arabidopsis MKS1 is involved with basal immunity and requires an intact N-terminal domain for proper function. [PLoS One 5; e14364](#).

Qiu JL, Fiil BK, Petersen K, Nielsen HB, Botanga CJ, Thorgrimsen S, Palma K, Suarez-Rodriguez MC, Sandbech-Clausen S, Lochota J, Brodersen P, Grasser KD, Mattsson O, Glazebrook J, Mundy J, Petersen M. 2008. Arabidopsis MAP kinase 4 regulates gene expression through transcription factor release in the nucleus. [EMBO J. 27; 2214-21](#).

Wiermer M, Cheng YT, Imkampe J, Li M, Wang D, Lipka V, Li X. 2012. Putative members of the Arabidopsis Nup107-160 nuclear pore sub-complex contribute to pathogen defense. [Plant Journal Jan 31](#).

Wiermer M, Germain H, Cheng YT, Garcia AV, Parker JE, Li X. 2010. Nucleoporin MOS7/Nup88 contributes to plant immunity and nuclear accumulation of defense regulators. [Nucleus 1\(4\); 332-336](#).

Wirthmueller L, Zhang Y, Jones JD, Parker JE. 2007. Nuclear accumulation of the Arabidopsis immune receptor RPS4 is necessary for triggering EDS1-dependent defense. [Curr Biol 17; 2023-9](#).

Yao C, Wu Y, Nie H, Tang D. 2012. RPN1a, a 26S proteasome subunit, is required for innate immunity in Arabidopsis. [Plant Journal, May 11](#).

Gary Lyon

December 2012